

## Use of Trees by Livestock: *ACACIA*

**J.J. Bennison and R.T.  
Paterson**

**Natural Resources  
Institute**

**32pp, 1993**

**ISBN: 0 85954 334-X**

---

➔ **Contents:**



**Foreword**



**Genus *Acacia***



**Summary**

**Description and distribution**



**Fodder characteristics**



**Anti-nutritive factors**



**Management**



**Alternative uses**



**References and further reading**



**Appendix 1 - Proximate and fibre analyses of bipinnate *Acacia* spp.**



**Appendix 2 - Proximate analyses of *Acacia* phyllodes**

---

© Crown copyright 1993

The Natural Resources Institute (NRI) is an internationally recognized centre of expertise on the natural resources sector in developing countries. It forms an integral part of the British Government's overseas aid programme. Its principal aim is to alleviate poverty and hardship in developing countries by increasing the

productivity of their renewable natural resources. NRI's main fields of expertise are resource assessment and farming systems, integrated pest management, food science and crop utilization.

NRI carries out research and surveys; develops pilot-scale plant, machinery and processes; identifies, prepares, manages and executes projects; provides advice and training; and publishes scientific and development material.

Short extracts of material from this publication may be reproduced in any non-advertising, nonprofit-making context provided that the source is acknowledged as follows:


Bennison, J.J., and Paterson, R.T. (1993) *Use of Trees by Livestock 2: Acacia*. Chatham, UK: Natural Resources Institute.

Permission for commercial reproduction should be sought from:

The Head, Publications and Publicity Section,  
Natural Resources Institute, Central Avenue,  
Chatham Maritime, Kent, ME4 4TB, United  
Kingdom.

No charge is made for single copies of this publication sent to the governmental and educational establishments, research institutions and non-profit-making organizations working in countries eligible for British Government Aid. Free copies cannot normally be addressed to individuals by name but only under their official titles.

Natural Resources Institute ISBN: 0 85954 334-X



## Use of Trees by Livestock: *ACACIA*



### **Foreword**



### **Genus *Acacia***



### **Summary**



### **Description and distribution**



### **Fodder characteristics**



### **Anti-nutritive factors**



### **Management**



### **Alternative uses**



### **References and further reading**



### **Appendix 1 - Proximate and fibre analyses of bipinnate *Acacia* spp.**



### **Appendix 2 - Proximate analyses of *Acacia* phyllodes**

---

## **Foreword**

**The importance of trees and shrubs in the feeding of animals in the tropics and sub-tropics has long been recognized by livestock owners. In arid areas where the growth of herbaceous plants is limited by lack of moisture, leaves and edible twigs of trees and shrubs can constitute well over 50% of the biomass production of rangeland. At high altitudes, tree foliage may provide over 50% of the feed available to ruminants in the dry season, branches being harvested and carried to the animals. Even in regions of higher rainfall where grass supplies the major proportion of the dry matter eaten by ruminants, tree leaves and fruits can form an important constituent of the diet, particularly for small ruminants.**


**In the last two decades interest in the planting of trees as a source of feed for livestock has been encouraged by workers in research and development, but in contrast to the hundreds of indigenous species which are used as fodder, attention has focussed on a limited number of introduced species. Thus**

**there are many publications reporting the chemical composition of *Leucaena leucocephala* leaves and suggesting management strategies for utilization of the tree for fodder, but it is more difficult to find information on alternative genera which might be equally, or more, appropriate.**

**The aim of this series of publications is to bring together published information on selected genera of trees which have the potential to increase the supply of fodder for ruminants. Each booklet summarizes published information on the fodder characteristics and nutritive value of one genus, with recommendations on management strategies, where available. Further, since the leaves of woody species frequently contain secondary compounds which may have an anti-nutritional, or toxic, effect, a separate booklet summarizes the effects of a number of these compounds. It is hoped that the booklets will provide useful resource material for students, research and extension workers, interested in promoting the use of trees as a source of fodder for ruminants**

**Further copies of this booklet or others in the series can be obtained by writing to the Publications Section at the Natural Resources Institute.**

**Margaret Gill**  
**Livestock Production Programme**



Use of Trees by Livestock: *ACACIA*



**Foreword**



**Genus *Acacia***



**Summary**



**Description and distribution**



**Fodder characteristics**



**Anti-nutritive factors**



**Management**



**Alternative uses**





## **References and further reading**



**Appendix 1 - Proximate and fibre analyses of bipinnate *Acacia* spp.**



**Appendix 2 - Proximate analyses of *Acacia* phyllodes**

---

## **Genus *Acacia***

**Family        LEGUMINOSAE**

**Subfamily   MIMOSOIDEAE**

**Tribe         ACACIEAE**

## **Principal phyllodenous species**

***Acacia aneura***

***Acacia combagei***

***Acacia coriacea***

***Acacia cyanophylla***

***Acacia dealbata***

***Acacia decurrens***

***Acacia holosericea***

***Acacia mearnsii***

***Acacia nilotica***

***Acacia pendula***

***Acacia salicina***

***Acacia tortilis***

***Acacia trachycarpa* (syn. *A. aff. linarioides*)**

***Acacia tumida***

## **Principal bipinnate species**

***Faidherbia albida* (syn. *Acacia albida*)**

***Acacia brevispica***

***Acacia cornigera***

***Acacia drepanolobium***

***Acacia farnesiana***

***Acacia haematoxylon* (syn. *A. giraffae*)**

***Acacia hebeclada* subsp. *hebeclada* (syn. *A. stolonifera*)**

***Acacia lasiopetala***

***Acacia leucophloea***

***Acacia mearnsii* (syn. *Racosperma mearnsii*)**

***Acacia mellifera***

***Acacia mellifera* subsp. *detinens***

***Acacia nilotica***

***Acacia oerfota* (syn. *A. nubica*)**

***Acacia robusta***

***Acacia senegal***

***Acacia seyal***

***Acacia sieberiana***


***Acacia tortilis***











***Acacia tortilis* subsp. *heteracantha* (syn. *A. likatunensis*)**



Use of Trees by Livestock: *ACACIA*

**Foreword**



-  **Genus *Acacia***
-  **Summary**
-  **Description and distribution**
-  **Fodder characteristics**
-  **Anti-nutritive factors**
-  **Management**
-  **Alternative uses**
-  **References and further reading**
-  **Appendix 1 - Proximate and fibre analyses of bipinnate *Acacia* spp.**
-  **Appendix 2 - Proximate analyses of *Acacia* phyllodes**

---

## Summary


**With the African *Acacia* spp. the fruit is probably more valuable as a feed than the foliage, because with the exception of *F. albida*, they tend to shed their leaves at the time of greatest feed scarcity during the dry season. Australian species however, retain their leaves and therefore show greater potential as fodder trees.**

**The productivity responses of animals reflect both the total intake, and the availability, of nutrients in the diet. Proximate and fibre analyses are unreliable indicators of nutritional value because of the presence of both cyanogenic glycosides and polyphenolics. A cautious approach should be adopted when feeding both *A. leucophloea* and some of the Australian species.**

**In some areas of Africa, *Acacia* spp. are the dominant browse trees within the natural ecosystem. Often there is no choice but to utilize the available feeds even if they do contain anti-nutritive compounds. Consumed as part of a wider diet, the**

**effect of anti-nutritive factors is diluted. They only become serious in periods of extreme feed scarcity or in situations where the opportunity for feed selection is reduced, such as in stall feeding and cut-and-carry systems.**

***Acacia* is a huge genus with pantropical distribution and a notable ability to survive in harsh environments. It has potential for use in environmental protection and for production of fodder, fuelwood, gums, and tannins.**



Use of Trees by Livestock: *ACACIA*



**Foreword**



**Genus *Acacia***



**Summary**



**Description and distribution**



**Fodder characteristics**



**Anti-nutritive factors**



**Management**



**Alternative uses**



**References and further reading**



**Appendix 1 - Proximate and fibre analyses of bipinnate *Acacia* spp.**



**Appendix 2 - Proximate analyses of *Acacia* phyllodes**

---

## **Description and distribution**

***Acacia* is a pantropical and subtropical genus with species abundant throughout Australia, Asia, Africa and the Americas. They thrive in a diverse range of habitats and environments. Many species are well adapted to the semi-arid and savannah regions but equally others survive in moist forest and riverine**

**areas, tolerating both high pH and waterlogged soils. With such diversity, *Acacia* has considerable potential in a range of livestock and agroforestry systems. In Africa and Australia, some naturally occurring species are important in traditional pastoral and agropastoral systems, while imported species have become commercially accepted. Trees provide fodder and shade for livestock, improve soil fertility through nitrogen fixation and the production of leaf litter and stabilize soils. *Acacia* species provide edible fruits and seeds, gum arabic and timber for fuel, construction and fencing.**

**In a large genus of 800-900 species (Allen and Allen, 1981), classification and nomenclature can be complex, particularly as there have been a number of revisions and attempts to subdivide the genus. Pedley (1986; 1987) advocated that it be divided into three separate genera, namely *Acacia*, with 200 species best represented in Africa and South America, *Senegalia*, with about 150 species with the same geographical distribution, and *Racosperma*, with about 850 species virtually**



**confined to Australia. While the suggestion may have merit, it has not yet been generally accepted. This text uses the traditional nomenclature adopted by standard flora. Where a name has been changed as a result of a generally accepted revision, the name used in the original reference is listed as a synonym. The names of the African species are those included in the check-list produced by Lock (1989) as part of the programme of the International Legume Database and Information Service (ILDIS).**

**One widely accepted recent change is the reclassification of *Acacia albida* as *Faidherbia albida*, although Coe and Beentje (1991) stated that the justification for the change was primarily phytochemical. In this publication, for purposes of discussion and comparison, the name *F. albida* is accepted but it is considered as though it had remained within the genus *Acacia*, since it shares many agronomic characteristics with species that have not been reclassified.**

**Various workers have divided the genus into divisions, series and subseries (Bentham, 1864) or sections (Taubert, 1894) but for ease of description and an understanding of the fodder potential, it is simplest to describe *Acacia* spp. as being either phyllodenous (principally Australian in origin) or bipinnate (mainly from Africa and Asia).**

### **Phyllodenous *Acacia* species**

**The majority of Australian *Acacia* spp. lack true leaves. Instead, they exhibit a characteristic known as phyllodony where the petiole is expanded and flattened. The phyllode has the ability to photosynthesize and for all practical purposes, it is the equivalent of a leaf. The Australian exceptions are from the Section *Botrycephalae*, which maintain bipinnate foliage until maturity and do not develop phyllodes.**

**Vercoe (1987) noted that species of *Acacia* and other genera from the dry tropics of Australia had potential in the semi-arid**


**zones of Africa for fuelwood and erosion control. If their usefulness as fodder could also be established, this would enhance their overall value. Certainly the value of trees such as *A. aneura* as drought reserve feed is well established and recognized by pastoralists in Australia. Unfortunately, success in establishing *A. aneura* in Africa has been limited. However, a number of other species appear to show potential, notably *A. holosericea*, *A. trachycarpa* and *A. coriacea*. Other species that have been studied include *A. salicina* and *A. cyanophylla*, although the phyllodes of the latter contain high tannin levels (Vercoe, 1987; Goodchild and McMeniman, 1987; Reed *et al.*, 1990), which can reduce both the palatability and the feeding value. Unlike African and Asian species, Australian *Acacia* spp. are evergreen and unarmed.**

### **Bipinnate *Acacia* species**

**There are about 200 bipinnate *Acacia* species in the New World and a further 150 species in Africa and Asia. All 129**

**African species possess bipinnate leaves. There are other physical and physiological characteristics which separate them from their Australian relatives, the *Botrycephalae*. This group, the best known of which is *A. mearnsii* (black wattle), grown throughout the subtropical world for the extraction of tannin, is confined to the cooler and moister parts of south eastern Australia. They are not noted as valuable fodder trees (Boland, 1987). Most bipinnate *Acacia* spp. are armed with thorns of various shapes and sizes, some of which are modified stipules at the base of the leaves. Many exhibit transient periods of defoliation in the dry season and have been described as deciduous. Before the start of the rains, there is often a flush of new growth of leaves. Although the stimulus for this activity is uncertain, it may be due to temperature.**





## Use of Trees by Livestock: *ACACIA*



### **Foreword**



### **Genus *Acacia***



### **Summary**



### **Description and distribution**



### **Fodder characteristics**



### **Anti-nutritive factors**



### **Management**



### **Alternative uses**



### **References and further reading**



### **Appendix 1 - Proximate and fibre analyses of bipinnate *Acacia* spp.**



### **Appendix 2 - Proximate analyses of *Acacia* phyllodes**

---

## **Fodder characteristics**

## Fodder yield

**Information on leaf yield of African bipinnate *Acacia* spp. is limited, and comparative data with introduced Australian species, even more so. The work of Cossalter (1986) compared species from the two regions, when grown in Senegal at uniform spacings of 3 x 3 m (Table 1). The results, as means of 20, or 25 individual trees aged between 18-23 months, demonstrated the fodder potential of the Australian *Acacia* species when compared with African ones. *A. holosericea* is perhaps the most frequently planted Australian *Acacia* in development projects because of its superior yield and because it retains a large phyllode biomass during the dry season, while African species shed their leaves during this period (Cossalter, 1986). There is some uncertainty about the palatability of *A. holosericea*: early trials indicate that only dry phyllodes are accepted. The fresh phyllodes of *A. trachycarpa* and *A. tumida*, however, were readily eaten by both cattle and**

**sheep (Hamel, 1980).****Table 1 Total biomass production (phyllodes, leaves and wood) of *Acacia* spp**

<b>Origin</b>	<b>Species</b>	<b>Dry leaves, phyllodes (g/tree)</b>	<b>Wood (kg/tree)</b>
<b>Australia</b>	<b><i>A. holosericea</i></b>	<b>2660</b>	<b>12.03</b>
<b>Australia</b>	<b><i>A. trachycarpa</i></b>	<b>1993</b>	<b>10.65</b>
<b>Australia</b>	<b><i>A. tumida</i></b>	<b>2600</b>	<b>8.69</b>
<b>Australia</b>	<b><i>A. trachycarpa</i></b>	<b>1700</b>	<b>3.73</b>
<b>Africa</b>	<b><i>A. senegal</i></b>	<b>60</b>	<b>9.92</b>
<b>Africa</b>	<b><i>A. seyal</i></b>	<b>208</b>	<b>9.39</b>
<b>Africa</b>	<b><i>A. tortilis</i></b>	<b>130</b>	<b>5.89</b>

**Source: Cossalter (1986)**

**When assessing the fodder value of African *Acacia* spp., the importance of the fruit and pods cannot be ignored. This fraction may be of more value than the foliage in some cases. A major problem in estimating the yield of fruit is the variation in individual trees between years. Trees may bear a heavy crop one year and little the next. The fruits of *F. albida* are relished by game, cattle and small ruminants and NAS (1979) suggests that yields of 125-135 kg are possible from a single tree. When ripe, the fruit can be harvested and stored for future use, although there is a tendency for the pods to shatter and disintegrate.**

**The bark of *A. seyal* is particularly valuable as a fodder, although complete removal will result in the death of the tree. Skerman *et al.*, (1988) suggested that the bark, which is extensively used for feeding cattle and small ruminants in the dry season, was the most valuable part of the plant. Cattle could consume up to 5.5 kg bark/day, sufficient for maintenance and the production of 4.5 litres of milk.**



## Nutritive value

**In Table 2, the mean proximate and fibre analyses of leaves, phyllodes and fruits of *Acacia* spp. are summarized from data of individual species (Appendix 1). Absolute values should be treated with caution since there is considerable variation in the literature. For example, the mean crude protein (CP) level of *Acacia* leaves is 18.9% of dry matter (DM), with a reported range of 6.5-42.8 %. However, the data provide an indication of the nutritive value and in general both leaves and fruits of all species are valuable sources of protein, particularly when compared with mature grasses, where CP levels in forage dry matter (DM) during the dry season can fall to 3-4% or less, well below the 7% CP which is suggested as a lower limit for fibre digestion in cattle (Van Soest, 1982).**

### **Table 2 Mean proximate and fibre analyses of *Acacia* leaves, phyllodes, fruits and seeds**

	<b>Crude protein</b>	<b>Crude fibre</b>	<b>Ash</b>	<b>Ether extract</b>	<b>NFE</b>	<b>NDF</b>	<b>ADF</b>
	(% of dry matter)						
<b><i>Acacia</i> LEAVES</b>							
<b>No. of data</b>	<b>41</b>	<b>41</b>	<b>40</b>	<b>39</b>	<b>38</b>		
<b>Mean</b>	<b>18.9</b>	<b>21.0</b>	<b>7.6</b>	<b>3.3</b>	<b>48.6</b>		
<b>Range</b>	<b>(7-43)</b>	<b>(8-46)</b>	<b>(5-12)</b>	<b>(2-13)</b>	<b>(32-66)</b>		
<b><i>Acacia</i> PHYLLODES</b>							
<b>No. of data</b>	<b>6</b>	<b>6</b>	<b>6</b>	<b>6</b>	<b>6</b>		
<b>Mean</b>	<b>12.2</b>	<b>27.6</b>	<b>7.0</b>	<b>2.9</b>	<b>48.7</b>		
<b>Range</b>	<b>(9-13)</b>	<b>(16-33)</b>	<b>(5-11)</b>	<b>(1-5)</b>	<b>(38-55)</b>		
<b><i>Acacia</i> FRUIT</b>							
<b>No. of data</b>	<b>33</b>	<b>29</b>	<b>33</b>	<b>28</b>	<b>28</b>	<b>4</b>	<b>4</b>

<b>Mean</b>	<b>14.5</b>	<b>24.7</b>	<b>5.8</b>	<b>1.7</b>	<b>51.9</b>	<b>34.6</b>	<b>25.7</b>
<b>Range</b>	<b>(8-22)</b>	<b>(12-41)</b>	<b>(3-14)</b>	<b>(1-3)</b>	<b>(31-67)</b>	<b>(32-37)</b>	<b>(23-28)</b>
<b>Acacia SEEDS</b>							
<b>No. of data</b>	<b>8</b>	<b>4</b>	<b>8</b>	<b>4</b>	<b>4</b>	<b>4</b>	<b>4</b>
<b>Mean</b>	<b>28.2</b>	<b>15.9</b>	<b>5.2</b>	<b>6.0</b>	<b>38.8</b>	<b>35.3</b>	<b>24.4</b>
<b>Range</b>	<b>(19-40)</b>	<b>(11-21)</b>	<b>(4-7)</b>	<b>(2-10)</b>	<b>(26-55)</b>	<b>(28-40)</b>	<b>(19-29)</b>

**Sources: Gohl, (1981); Goodchild and McMeniman, (1987); Skerman *et al.*, (1988); Tanner *et al.*, (1990)**

**Notes: NFE - Nitrogen-free extract; NDF - Neutral detergent fibre; ADF - Acid detergent fibre.**

**While the crude fibre (CF) levels of *Acacia* are relatively high (15-40%) compared with other fodder tree species such as *Leucaena leucocephala* and *Gliricidia sepium*, they are still markedly lower than for tropical grasses and common crop**

**residues. Even the fruit, consisting of the dry dehiscent pod and the seeds contained in it, are a valuable feed source, with moderate CF (25%) and high CP levels (15%). The phyllodenous *Acacia* spp. tend to have marginally higher CF and lower CP levels than bipinnate species (Appendix 2) though the available data consist of fewer observations. The lower nutritive value is offset by both the higher biomass yield and the retention of the phyllodes throughout the dry season, when bipinnate species tend to shed their leaves.**

**Information on the digestibility of DM and organic matter (OM) of *Acacia* leaves and phyllodes is limited but the available data indicate that, with sheep, they are relatively low and in the range of 40-60% respectively (Table 3). The digestibility of phyllodes appears to be lower than that of bipinnate leaves. Goodchild and McMeniman, (1987) suggested that the low fibre digestibility could be associated with the high lignin content of the cell wall, since fibre digestibility is inversely related to the lignin content of the**

## **fibre fraction.**

**While the digestibility of the whole fruit (seed and pod) is higher than that of the leaves and phyllodes, it can be adversely affected by the proportion of seeds that pass undigested through the gut of the animal. Tanner *et al.*, (1990) reported that 46% of *A. tortilis* seeds were voided in the faeces of sheep. Similarly, Goodchild and McMeniman (1987) noted that when feeding yearling cattle with seeds of *A. tortilis* subsp. *spirocarpa*, 12% passed unchanged through the gut. Legume seeds are physically hard and notoriously resistant to digestive juices, although the small size of *Acacia* seed may be a contributory factor in their passage through the gut (Tanner *et al.*, 1990).**

**The digestibility of the leaf and fruit fraction can also be depressed by the presence of polyphenolic compounds, including tannins.**


**Table 3 *In vivo* digestibility (%) of *Acacia* forage**

	<b>Organic matter</b>	<b>Dry matter</b>	<b>Neutral detergent fibre</b>	<b>Nitrogen</b>	<b>Crude protein</b>	<b>Source</b>
<b>LEAVES</b>						
<i>A. farnesiana</i>	<b>60*</b>				<b>19.8</b>	<b>1</b>
<i>F. albida</i>	<b>53</b>				<b>17.8</b>	<b>1</b>
<i>A. sieberiana</i>	<b>54</b>		<b>37</b>	<b>27</b>	<b>16.9</b>	<b>3</b>
<i>A. seyal</i>	<b>54</b>		<b>42</b>	<b>28</b>	<b>18.2</b>	<b>3</b>
<b>PHYLLODES</b>						
<i>A. aneura</i>	<b>47</b>	<b>32*</b>			<b>12.2</b>	<b>1,4</b>
<i>A. combagei</i>	<b>44*</b>					<b>1</b>
<i>A. pendula</i>	<b>43</b>					<b>1</b>

<b>A. holosericea</b>		<b>32*</b>				<b>4</b>
<b>A. cyanophylla</b>	<b>41</b>		<b>29</b>	<b>10</b>	<b>6.3</b>	<b>3</b>
<b>FRUIT (pods and seeds)</b>						
<b>A. tortilis</b>	<b>61</b>	<b>58</b>	<b>51</b>	<b>30</b>	<b>14.5</b>	<b>2</b>
<b>F. albida</b>	<b>57</b>	<b>55</b>	<b>45</b>	<b>29</b>		<b>2</b>
<b>F albida</b>	<b>58</b>	<b>1</b>			<b>11.0</b>	<b>1</b>
<b>A nilotica</b>	<b>61</b>	<b>59</b>	<b>49</b>	<b>32</b>		<b>2</b>
<b>A. nilotica</b>	<b>66</b>	<b>1</b>		<b>29</b>	<b>11.3</b>	
<b>A. sieberiana</b>	<b>55</b>	<b>53</b>	<b>39</b>		<b>18.1</b>	<b>2</b>

**Notes: \* *in vitro* determinations.**

**Sources: 1 Goodchild and McMeniman (1987); 2 Tanner *et al.* (1990); 3 Reed *et al.* (1990); 4 Vercoe (1987).**



Use of Trees by Livestock: *ACACIA*



**Foreword**



**Genus *Acacia***



**Summary**



**Description and distribution**



**Fodder characteristics**



**Anti-nutritive factors**



**Management**



**Alternative uses**



**References and further reading**





**Appendix 1 - Proximate and fibre analyses of bipinnate *Acacia* spp.**



**Appendix 2 - Proximate analyses of *Acacia* phyllodes**

---

## **Anti-nutritive factors**

***Acacia* spp. have acquired a variety of physical and chemical defence mechanisms aimed at reducing the palatability and nutritive value of the plant to predators including both insects and animals.**

**Perhaps the most obvious physical characteristic is the spiny armature of the African species, which evolved as a defence against large herbivores. *A. mellifera* subsp. *detinens* derived its common name (wait-a-bit tree) from the pairs of small hooks on the branches. *A. tortilis* is protected by two types of thorns, long white stipular spines and small curved hooks, a formidable protection against animals. Coe and Beentje**

**(1991) suggested that as thorns restrict the availability of forage, a critical level is reached below which the herbivore no longer finds it nutritionally efficient to seek shoots between the spines, since this invariably involves movement between each bite. There is also evidence that in *A. drepanolobium*, thorns on branches which are regularly browsed by animals are significantly longer than those on undefoliated branches (Young, 1987).**

**Tree size is a form of physical protection that can be harnessed as a desirable characteristic in certain agroforestry systems. Domestic animals are unable to browse tall trees which then maintain a standing reserve of foliage which can be made available by strategic lopping at critical periods when other feed is scarce.**

**A fascinating defence mechanism of the genus is the symbiotic relationship with ants known as myrmecophily, where the plant lives in association with an ant colony. In the tropical**

**American species *A. cornigera*, association with ants from the genus *Pseudomyrmex* is a notable example of myrmecophily. The trees have modified, swollen spines or galls, which provide shelter for the ants, and extrafloral nectaries and modified leaflet tips which provide food. The ants, in return, protect the trees from insects and other predators by swarming to attack them when the tree is disturbed (Janzen, 1966). In Africa, *A. drepanolobium* is commonly known as the 'whistling thorn', a name derived from the noise of the wind blowing into the holes of the galls that have been occupied by ants.**

**Many *Acacia* spp. produce a range of potentially toxic compounds which prevent herbivores from eating the foliage, although it is thought that the chemicals originally evolved as defence mechanisms against insect and fungal attack. Some members contain specific anti-nutritive factors, for example, *A. georginae* has been implicated in heavy stock losses in Australia due to its content of fluoracetic acid (D'Mello, 1992).**

**In general terms the genus contains two major groups of anti-nutritive factors, cyanogenic glycosides and polyphenolic compounds.**

## **Cyanogenic glycosides**

**When cyanogenic substrates are broken down under the action of a hydrolyzing enzyme, hydrogen cyanide (HCN) is formed. Plants that contain both cyanogenic glycosides and an endogenous, hydrolytic enzyme are the most dangerous but broad-spectrum enzymes may be present in other fractions of the diet. Symptoms of HCN poisoning include respiratory dyspnoea, intense red conjunctiva, frothing at the mouth, bloat, a staggering gait, convulsions and violent death. On post-mortem examination, a characteristic almond smell is often noted in the contents of the gut.**

**In Asia, there are a number of reports of HCN poisoning of livestock when fed on the pods and leaves of *A. leucophloea***

**(Gupta and Naurival, 1966; Prasad *et al.*, 1977; Vihan and Panwar, 1987). In South Africa, Steyn and Rimington (1935) reported the presence of cyanogenic glycosides in *A. haematoxilon*, *A. lasiopetala*, *A. tortilis* subsp. *heteracantha*, *A. robusta* and *A. hebeclada* subsp. *hebeclada*. They found higher concentrations of HCN from leaves and immature pods than from mature pods. In *A. robusta*, the glycoside was present without the corresponding hydrolytic enzyme. They isolated the cyanogenic compound acacipetalin from *A. lasiopetala* and *A. hebeclada*. Siegler *et al.*, (1975) isolated a further compound, dihydroacipetalin, from the leaves and shoots of *A. sieberiana* var. *woodii*.**

**Toxicity appears only in sporadic outbreaks, being dependant on a number of factors, including the species and variety of *Acacia*, the rate of ingestion by the animal and the availability of alternative feeds. It is more likely to occur during periods of drought or feed scarcity when hungry animals consume large quantities of feed in a short period of time. Access to**

**cold water during feeding appears to promote the release of HCN from the substrates. Practical measures can be taken to reduce the risk of poisoning such as restricting the intake, particularly of leaves and immature pods, separating the time or place of feeding and watering, and feeding dangerous material with molasses or sulphur. These substances combine with HCN to form harmless cyanhydrin and sulphocyanides respectively (Steyn and Rimmington, 1935).**

## **Phenolic compounds**

**There are numerous reports of the presence of polyphenolic compounds in the various components of plants of the genus *Acacia*. From the standpoint of nutrition, the most important of these are the tannins, compounds with molecular weights in excess of 500, which are able to form complexes with, and to precipitate proteins (McLeod, 1974; Mangan, 1988). Hydrolyzable tannins are readily broken down by acids, bases and enzymes to form sugars and phenolic carboxylic acid and**

**are therefore easily degraded in the gut. In contrast, condensed tannins are not degraded and so are available to form complexes with proteins, resulting in reduced feed intake and digestibility. McNaughton (1987) suggested that at levels above about 5%, increasing the tannin content in a diet reduced its palatability.**

**Reed *et al.*, (1990) assessed the effects of polyphenolics on intake, growth, digestibility and nitrogen utilization by sheep. Three *Acacia* forages, (phyllodes of *A. cyanophylla*, leaves of *A. seyal*, and fruit of *A. sieberiana*), all containing high levels of polyphenolics, were compared with three protein supplements and another fodder tree *Sesbania sesban* which is low in polyphenolics. All diets were fed in combination with teff straw (*Eragrostis abyssinica*). The diets containing *Acacia* led to the highest levels of feed refusal by the sheep their and lowest intake of teff straw. Fibre digestibility was lower for these diets and the tannins complexed with proteins, resulting in higher levels of faecal nitrogen.**


**A complementary study by Tanner *et al.*, (1990) reported significant differences in the growth rates of sheep supplemented with fruits from four different *Acacia* spp. (*A. tortilis*, *F. albida*, *A. nilotica* and *A. sieberiana*) even though there were no significant differences in total nitrogen intake. The tannin-rich fruits, particularly of *A. sieberiana* and *A. nilotica*, reduced feed intake and the efficiency of nitrogen utilization by forming complexes with proteins which again resulted in a shift of nitrogen excretion from urine to faeces. The growth rates of sheep fed the diets low in polyphenolics (*F. albida* and *A. tortilis*), were not significantly different from sheep supplemented with noug meal (*Guizotia abyssinica*), indicating their suitability as dietary supplements.**

**The literature abounds with conflicting reports on the suitability of *Acacia* spp. for livestock feed. Dumancic and Le Houerou (1981) considered the phyllodes of *A. cyanophylla* to be sufficient to meet the protein maintenance requirements of sheep, while Reed *et al.*, (1990) reported a negative nitrogen**



**balance for animals fed on this legume. Predictions of the productive responses of livestock are problematical and uncertain. There are a number of possible explanations for the conflicting reports:**

- **the chemistry of tannins is complex and analytical results do not always correlate with biological effects, e.g. nitrogen digestibility;**
- **a single feed may contain more than one antinutritive factor. *A. nilotica* is known to have both tannins and catechin gallates (Mueller-Harvey et al., 1987);**
- **the concentration and type of phenolic compound both vary with season and in response to stimuli such as defoliation;**
- **there is much variation between both species and provenance in tannin content, the effect being both genetic and environmental.**



## Use of Trees by Livestock: *ACACIA*



### **Foreword**



### **Genus *Acacia***



### **Summary**



### **Description and distribution**



### **Fodder characteristics**



### **Anti-nutritive factors**



### **Management**



### **Alternative uses**



### **References and further reading**



### **Appendix 1 - Proximate and fibre analyses of bipinnate *Acacia* spp.**



### **Appendix 2 - Proximate analyses of *Acacia* phyllodes**

## **Management**

**Where the principal use of *Acacia* spp. is for browsing by livestock, little is done to manage the resource, except through some measure of defoliation control by herding the animals. High branches are lopped during times of feed shortage, particularly in the dry season. At other times of the year, the trees are subjected to uncontrolled browsing, often by both wild and domestic animals.**

**Many legumes produce seed which has a hard, impermeable seed coat. Such seed will not germinate until moisture is able to penetrate it. This characteristic, known as 'hard seededness', prevents the simultaneous germination of the entire stock of seed in the soil at the start of the wet season. This adaptation favours the long-term survival of the population where rainfall is unpredictable. It is, however, a disadvantage when attempting to establish new stands from seed, particularly in areas with a short growing season. *Acacia***

**shows extreme variability in the degree of hard seededness, both between and within species; the proportion of hard seed in a sample depends on the environmental conditions during plant growth, the maturity of the seed at harvest and the length of time it has spent in storage. In general, Australian species respond to seed scarification. Recommended treatments (Doran and Gunn, 1987) include:**


- **manual nicking of the seed coat. Because of the amount of labour involved, this is suitable only for seed for germination tests, or for valuable research lots;**
- **placing the seed in boiling water for one minute. This is recommended for many hard-coated seed lots;**
- **placing the seed in hot water (80-90°C) for one minute. This is recommended where sensitivity to boiling water is suspected.**

**In Central Australia where annual rainfall is about 300 mm, native *Acacia* spp. naturally regenerate after exceptionally**

**high rainfall at populations of some 5000 stems/ha, and many species produce multiple stems. Subsequent fires and drought will thin the stand to 100-400 stems/ha. Under these conditions, the optimum stocking level for planted stands may be in the region of 1000 seedlings/ha, followed by thinning as the stand matures. Alternatively, where thinning is not practical, an initial stand of about 400 trees/ha would be appropriate (Kube,1987). The seed is scarified in hot water (80°C) and germinated in trays. Seedlings are transplanted after a few days into individual tubes 5 cm in diameter by 15 cm deep, filled with a potting mixture of equal parts of coarse sand, fine sand or sandy loam and peat moss. Treatment with fungicide is usually necessary and the seedlings are grown in decreasing shade for about three months before planting out. Supplementary irrigation is applied for 6-12 months, depending upon rainfall. Growth is often rapid, many species attaining heights of 2 m in about two years.**

**In Tanzania, *A. mearnsii* plantations have been established for**

**fuelwood, charcoal and building poles by direct seeding scarified seed at rates of 1.2 - 2.4 kg/ha in rows 1.8 m apart in fully cultivated soil (Kessy,1987). The major problem with this species has been its susceptibility to competition from weeds. Harvesting has been on an 8-12 year cycle and natural regeneration has taken place after the original stand has been clear felled and burned.**



Use of Trees by Livestock: *ACACIA*



**Foreword**



**Genus *Acacia***









**Summary**



**Description and distribution**



**Fodder characteristics**

-  **Anti-nutritive factors**
-  **Management**
-  **Alternative uses**
-  **References and further reading**
-  **Appendix 1 - Proximate and fibre analyses of bipinnate *Acacia* spp.**
-  **Appendix 2 - Proximate analyses of *Acacia* phyllodes**

---

## Alternative uses

**While *Acacia* is the largest of the mimosoid genera and the second largest of all the legume family, only about 75 members have economic value, and of these only about 50 are actually cultivated. Nevertheless, they are used in various ways, as described by Allen and Allen, (1981) and Boland, (1987).**

**Many *Acacia* spp. form spiny, slow-growing scrub which provides protection against soil erosion caused by wind and rain. Their tolerance of harsh conditions makes them valuable for stabilization and revegetation of difficult sites. *A. armata*, *A. glaucescens*, *A. pycnantha* and *A. suaveolens* tolerate salt spray and so can be used to stabilize coastal sand dunes. *A. decurrens* is planted along roadsides in Australia and on denuded hills in Rwanda. *A. mearnsii* and *A. dealbata* are used to control gulley and hillside erosion in New Zealand.**

***Acacia* is an important source of fuelwood or charcoal in arid regions. *A. mearnsii* is used for both domestic fuel and tobacco curing in Indonesia (NAS, 1980). It also provides poles for mining and agriculture in many parts of the world.**

**As timber, *Acacia* wood is extremely durable and although it is hard to work, it finishes well. Smaller species provide wood which is used to make implement handles, pipes and furniture, while larger species are used for panelling, boats and musical**




**instruments. Hawaiian mahogany (*A. koa*) was once prized for making surfboards, canoes and bowls while *A. segal* and *A. tortilis* were the principal trees of the Arabian Desert region. *A. glaucescens* and *A. melanoxylon* are major timber trees today. *A. dealbata* and *A. mearnsii* are used for pulping in Australia and Japan, while the latter species is used in the production of rayon in South Africa and India.**






***A. mearnsii* is the world's most important source of vegetable tannin, 35-40% of the air-dry weight of bark is extracted for leather manufacture. Yields in excess of 15% are also obtainable from other species such as *A. albida* and *A. cyanophylla*. Wood adhesives and polyurethane foams can be prepared from bark extracts of *A. mearnsii*, and used as substitutes for petroleum products.**









**Gum arabic, an important emulsifier, protective colloid, adhesive and binder, is produced in commercially exploitable quantities by *A. arabica*, the principal historical source, and**

more recently by *A. senegal* and some 20 other species of *Acacia*. It is used in foods, pharmaceuticals, cosmetics, adhesives, paints, polishes and inks and in lithography, photography, textile-sizing and other processes.

Species such as *A. dealbata*, known as mimosa by florists, *A. albida* and *A. tortilis*, are valued as ornamentals while *A. farnesiana* and *A. dealbata* are used in the production of base oils for perfumes. In Australia, several species are valuable sources of pollen in apiculture.



-  Use of Trees by Livestock: *ACACIA*
-  **Foreword**
-  **Genus *Acacia***
-  **Summary**
-  **Description and distribution**

-  **Fodder characteristics**
-  **Anti-nutritive factors**
-  **Management**
-  **Alternative uses**
-   **References and further reading**
-  **Appendix 1 - Proximate and fibre analyses of bipinnate *Acacia* spp.**
-  **Appendix 2 - Proximate analyses of *Acacia* phyllodes**

---

## References and further reading

**AHN, J.H., ROBERTSON, B.M., ELLIOT, R., GUTTERIDGE, R.C. and FORD, C.W. (1989) Quality assessment of tropical browse legumes: tannin content and protein degradation. *Animal Feed Science and Technology*, 27: 147-156**

**ALLEN, O.N. and ALLEN, E.K. (1981) *The Leguminosae: a Source Book of Characteristics, Uses and Nodulation*. Madison, USA: University of Wisconsin Press.**

**BATE-SMITH, E.C. (1973) Haemanalysis of Tannins: The Concept of Relative Astringency. *Phytochemistry*, 12: 907-912.**

**BENTHAM, G. (1864) *Flora Australiensis* vo1.2. London: Reeve and Co.**

**BOLAND, D. J. (1987) Genetic resources and utilisation of Australian bipinnate acacias (Botrycephalae). In: *Australian Acacias in Developing Countries*. TURNBULL, J.W. (ed.) Proceedings, International Workshop held at Gympie, Australia, August, 1986. ACIAR Proceedings No. 16. pp. 29-37.**

**BUTLER, L.G. (1982) Relative degree of polymerization of sorghum tannin during seed development and maturation. *Journal of Agriculture and Food Chemistry*, 30:1090-1094.**

**COE, M. and BEENTJE, H. (1991) *A Field Guide to the Acacias of Kenya*. Oxford, UK: Oxford University Press.**

**COSSALTER, C. (1986) Introduction of Australian Acacias into Dry Tropical West Africa. *Forest Ecology and Management*, 16: 367-389.**

**D'MELLO, J.P.F. (1992) Chemical Constraints to the use of tropical legumes in animal nutrition. *Animal Feed Science and Technology*, 38: 237-261**

**DORAN, J.C. and GUNN, B.V. (1987) Treatments to promote seed germination in Australian Acacias. In: *Australian Acacias in Developing countries*. TURNBULL, J.W. (ed.) Proceedings, International Workshop held at Gympie, Australia, August 1986. ACIAR Proceedings No. 16. pp. 57-63**

**DUMANCIC, D. and LE HOUEROU, H.N. (1981) *Acacia***

***cyanophylla* Lindl. as a supplementary feed for small stock in Libya. *Journal of Arid Environments*, 4:161-167.**

**GOHL, B. (1981) *Tropical Feeds: Feed Information Summaries and Nutritive Values*. FAO Animal Production and Health Series No. 12. Rome: FAO.**

**GOODCHILD, AN. and McMENIMAN, N.P. (1987) Nutritive value of *Acacia* foliage and pods for animal feeding. In: *Australian Acacias in Developing Countries*. TURNBULL, J.W. (ed.) Proceedings, International Workshop held at Gympie, Australia, August 1986. ACIAR Proceedings No.16. pp. 101-106.**

**GUPTA, I. and NAURIYAL, M.M. (1966) *Acacia leucophloea* Willd (raunja) poisoning in livestock. *Indian Veterinary Journal*, 43: 538-543.**

**HAGERMAN, A.E. and BUTLER, L.G. (1981) The specificity of**

**proanthocyanidin-protein interactions. *Journal of Biological Chemistry*, 256: 4494-4497.**

**HAMEL, O. (1980) Acclimatization and utilization of phyllodineous acacias from Australia in Senegal. In: *Browse in Africa; The current state of knowledge*. LE HOUEROU, H.N. (ed.) International Livestock Centre for Africa: Addis Ababa. pp.361-373.**

**JANZEN, D.H. (1966) Coevolution of mutualism between ants and acacias. *Evolution*, 20: 249-275.**

**KESSY, B.S. (1987) Growth of Australian Acacias in Tanzania. In: *Australian Acacias in Developing Countries*. TURNBULL, J.W. (ed.) Proceedings, International Workshop held at Gympie, Australia, August 1986. ACIAR Proceedings No. 16. pp. 123-125.**

**KUBE, P.D. (1987) Growth rates, establishment techniques**

**and propagation of some Central Australian Acacias. In: *Australian Acacias in Developing Countries*. TURNBULL, J.W. (ed.) Proceedings, International Workshop held at Gympie, Australia, August 1986. ACIAR Proceedings No. 16. pp. 77-80.**

**KUMAR, R. and VAITHIYANATHAN, S. (1990) Occurrence, nutritional significance and effect on animal productivity of tannins in tree leaves. *Animal Feed Science and Technology*, 30: 21-38.**

**LE HOUEROU, H.N. (1980) The role of browse in the Sahelian and Sudanian zones. In: *Browse in Africa; The current state of knowledge*. LE HOUEROU, H.N. (ed.) International Livestock Centre for Africa: Addis Ababa. pp.83-100.**

**LOCK, J.M. (1989) *Legumes of Africa: A Check List*. Kew: Royal Botanic Gardens.**

**MAKKAR, H.P.S., DAWRA, R.K. and SINGH, B. (1988) Changes**



**in tannin content, polymerisation and protein precipitation capacity in oak (*Quercus incana*) leaves with maturity. *Journal of Science of Food and Agriculture*, 44: 301-307.**

**MANGAN, J.L. (1988) Nutritional effects of tannins in animal feeds. *Nutrition Research Reviews*, 1:209-231.**

**MASLIN, B.R., CONN, E.E. and DUNN, J.E. (1987) Cyanogenic Australian species of *Acacia*: A preliminary account of their toxicity potential. In: *Australian Acacias in Developing Countries*. TURNBULL, J.W. (ed.) Proceedings, International Workshop held at Gympie, Australia, August 1986. ACIAR Proceedings No. 16. pp. 107-111.**

**McLEOD, M.N. (1974) Plant tannins - their role in forage quality. *Nutrition Abstracts and Reviews*, 44: 803-815.**

**McNAUGHTON, S.J. (1987) Adaption of herbivores to season changes in nutrient supply. In: *Nutrition of Herbivores*.**

**HACKER, J.B. and TERNMOUTH, J.H. (eds). Sydney, Australia: Academic Press. pp. 391-408.**

**MUELLER-HARVEY, I., REED, J.D. and HARTLEY, R.D. (1987) Characterisation of phenolic compounds, including flavonoids and tannins, of ten Ethiopian browse species by high performance liquid chromatography. *Journal of the Science of Food and Agriculture*, 39: 1-14.**

**NAS (1979) *Tropical Legumes: Resources for the Future*. Washington DC: National Academy of Sciences.**

**NAS (1980) *Firewood Crops*. Washington DC: National Academy of Sciences.**

**PEDLEY, L. (1986) Deprivation and dispersal of *Acacia* (*Leguminosae*) with particular reference to Australia, and recognizing *Senegalia* and *Racosperma*. *Botanical Journal of the Linnean Society*, 92: 219-54.**

**PEDLEY, L. (1987) Australian Acacias: Taxonomy and Phytogeography. In: *Australian Acacias in Developing Countries*. TURNBULL, J.W.(ed.) Proceedings, International Workshop held at Gympie, Australia, August 1986. ACIAR Proceedings No 16. pp. 11-16.**

**PRASAD, J., SINGH, A.P. and REKIB, A. (1977) Hydrocyanic acid poisoning in grazing sheep and goat on *Acacia leucophloea* (reunja). *Indian Veterinary Journal*, 54:748-751.**

**REED, J.D., HORVARTH, P.J., ALLEN M.S. and VAN SOEST, P.J. (1985) Gravimetric determination of soluble phenolics including tannins from leaves by precipitation with trivalent ytterbium. *Journal of the Science of Food and Agriculture*, 36: 255-261.**

**REED, J.D., SOLLER, H., and WOODWARD, A. (1990). Fodder tree and straw diets for sheep: intake, growth, digestibility**

**and the effects of phenolics on nitrogen utilisation. *Animal Feed Science and Technology*, 30: 3950.**

**SEDDON, H.R. and KING, R.O.C. (1930) As to the fatal dose for sheep of cyanogenetic plants containing Sambunigrin and Prunasin. *Journal Council Scientific and Industrial Research Australia*, 3: 14-24.**

**SIEGLER, D.S, BUTTERFIELD, C.S., DUNN, J.E. and CONN, E.E. (1975). Dihydroacacipetalin - a new cyanogenic glycoside from *Acacia sieberiana* var *woodii*. *Phytochemistry*, 14: 1419-1420.**

**SKERMAN, P.J., CAMERON, D.G. and RIVEROS, F. (1988) *Tropical Forage Legumes*. 2nd. edn. Rome: FAO.**

**STEYN, D.G. and RIMINGTON, C. (1935). The occurrence of cyanogenetic glucosides in South African species of *Acacia*. *Onderstepoort Journal of Veterinary Science and Animal***

***Industry, 4(1): 51-73.***

**TANNER, J.C., REED, J.D. and OWEN, E. (1990). The nutritive value of fruits (pods with seeds) from four *Acacia* spp. compared with extracted noug (*Guizotia abyssinica*) meal as supplements to maize stover for Ethiopian highland sheep. *Animal Production, 51:127-133.***

**TAUBERT, P. (1894) *Die Naturlichen Pflanzenfamilie*. Leipzig: Wilhelm Engelmann.**

**TERBLANCE, M., PIENAAR, J.G., BIGALKE, R. and VAHRMEYER, J. (1967) *Acacia nilotica* (L.) Del. subsp. *kraussiana* (Benth) Brenan. as a poisonous plant in South Africa. *Journal South African Veterinary Medical Association, 38(1): 57-63.***

**VAN HOVEN, W. (1985) The tree's secret weapon. *South African Panorama, 30(3): 34-37.***

**VAN SOEST, P.J. (1982) *Nutritional Ecology of the Ruminant*. Corvallis, Oregon, USA: O & B Books.**

**VERCOE, T.K. (1987) Fodder potential of selected Australian tree species. In: *Australian Acacias in Developing Countries*. TURNBULL, J.W. (ed.) Proceedings, International Workshop held at Gympie, Australia, August 1986. ACIAR Proceedings No. 16. pp. 95-100.**


**VIHAN, V.S. and PANWAR, H.S. (1987) A note on toxicity of *Acacia leucophloea* (Rajma) in sheep. *Indian Journal Animal Research*, 21(1): 53-55.**









**WILSON, A.D. (1969). A review of browse in the nutrition of grazing animals. *Journal of Range Management*, 22:23-28.**

**WOODWARD, A. and REED, J.D. (1989) The influence of polyphenolics on the nutritive value of browse: A summary of research conducted at ILCA. *ILCA Bulletin 35*, Addis Ababa:**

## International Livestock Centre for Africa.

**YOUNG, T.P. (1987) Increased thorn length in *Acacia drepanolobium* - an induced response to browsing. *Oecologia* 71(3): 436-438.**



-  Use of Trees by Livestock: *ACACIA*
-  **Foreword**
-  **Genus *Acacia***
-  **Summary**
-  **Description and distribution**
-  **Fodder characteristics**
-  **Anti-nutritive factors**
-  **Management**

**Alternative uses****References and further reading****Appendix 1 - Proximate and fibre analyses of bipinnate *Acacia* spp.****Appendix 2 - Proximate analyses of *Acacia* phyllodes**

---

**Appendix 1 Proximate and fibre analyses of bipinnate *Acacia* spp.**

	Dry matter %	Crude protein (% of dry matter)	Crude fibre	Ash	Ether extract	NFE	NDF	ADF
<b><i>Faidherbia albida</i> LEAVES</b>								



<b>No. of data</b>	<b>1</b>	<b>4</b>	<b>4</b>	<b>4</b>	<b>3</b>	<b>2</b>	<b>0</b>	<b>0</b>
<b>Low</b>	<b>36.3</b>	<b>17.1</b>	<b>12.4</b>	<b>6.4</b>	<b>1.6</b>	<b>51.9</b>		
<b>High</b>	<b>36.3</b>	<b>19.7</b>	<b>19.6</b>	<b>8.4</b>	<b>3.7</b>	<b>59.8</b>		
<b>Mean</b>	<b>36.3</b>	<b>18.1</b>	<b>16.8</b>	<b>7.4</b>	<b>2.5</b>	<b>55.8</b>		
<b><i>Faidherbia albida</i> FRUIT</b>								
<b>No. of data</b>	<b>0</b>	<b>6</b>	<b>5</b>	<b>6</b>	<b>4</b>	<b>4</b>	<b>1</b>	<b>1</b>
<b>Low</b>		<b>8.8</b>	<b>21.4</b>	<b>2.7</b>	<b>0.7</b>	<b>42.6</b>	<b>37.4</b>	<b>27.9</b>
<b>High</b>		<b>14.3</b>	<b>40.7</b>	<b>6.3</b>	<b>1.5</b>	<b>61.7</b>	<b>37.4</b>	<b>27.9</b>
<b>Mean</b>		<b>12.0</b>	<b>27.8</b>	<b>4.0</b>	<b>1.1</b>	<b>53.6</b>	<b>37.4</b>	<b>27.9</b>
<b><i>Faidherbia albida</i> PODS</b>								
<b>No. of data</b>	<b>0</b>	<b>1</b>	<b>0</b>	<b>1</b>	<b>0</b>	<b>0</b>	<b>1</b>	<b>1</b>
<b>Low</b>		<b>7.8</b>		<b>3.8</b>			<b>44.3</b>	<b>31.9</b>
<b>High</b>		<b>7.8</b>		<b>3.8</b>			<b>44.3</b>	<b>31.9</b>
<b>Mean</b>		<b>7.8</b>		<b>3.8</b>			<b>44.3</b>	<b>31.9</b>

<b><i>Faidherbia albida</i> SEEDS</b>								
<b>No. of data</b>	<b>0</b>	<b>1</b>	<b>0</b>	<b>1</b>	<b>0</b>	<b>0</b>	<b>1</b>	<b>1</b>
<b>Mean</b>		<b>27.9</b>		<b>4.9</b>			<b>29.7</b>	<b>18.2</b>
<b><i>Acacia brevispica</i> LEAVES</b>								
<b>No. of data</b>	<b>0</b>	<b>5</b>	<b>5</b>	<b>5</b>	<b>5</b>	<b>5</b>	<b>0</b>	<b>0</b>
<b>Low</b>		<b>17.1</b>	<b>18.0</b>	<b>5.6</b>	<b>2.0</b>	<b>41.5</b>		
<b>High</b>		<b>26.0</b>	<b>29.9</b>	<b>8.2</b>	<b>5.7</b>	<b>51.0</b>		
<b>Mean</b>		<b>19.6</b>	<b>23.5</b>	<b>6.6</b>	<b>3.7</b>	<b>46.7</b>		
<b><i>Acacia drepanolobium</i> LEAVES</b>								
<b>No. of data</b>		<b>1</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>1</b>		
<b>Mean</b>		<b>15.4</b>	<b>29.8</b>	<b>5.5</b>	<b>2.3</b>	<b>47.0</b>		

<b><i>Acacia farnesiana</i></b> <b>LEAVES</b>								
<b>No. of data</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>1</b>		
<b>Mean</b>	<b>40.0</b>	<b>17.2</b>	<b>17.9</b>	<b>5.2</b>	<b>2.1</b>	<b>57.1</b>		
<b><i>Acacia farnesiana</i></b> <b>FRUIT</b>								
<b>No. of data</b>	<b>0</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>1</b>		
<b>Mean</b>		<b>17.2</b>	<b>19.4</b>	<b>4.1</b>	<b>1.6</b>	<b>57.7</b>		
<b><i>Acacia farnesiana</i></b> <b>SEEDS</b>								
<b>No. of data</b>		<b>1</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>1</b>		
<b>Mean</b>		<b>20.9</b>	<b>18.3</b>	<b>3.8</b>	<b>2.3</b>	<b>54.7</b>		
<b><i>Acacia</i></b>								

<b><i>mellifera</i></b> <b>LEAVES</b>								
<b>No. of data</b>	<b>1</b>	<b>4</b>	<b>4</b>	<b>4</b>	<b>4</b>	<b>4</b>		
<b>Low</b>	<b>34.1</b>	<b>8.0</b>	<b>14.1</b>	<b>6.2</b>	<b>1.8</b>	<b>32.2</b>		
<b>High</b>	<b>34.1</b>	<b>42.8</b>	<b>45.8</b>	<b>10.2</b>	<b>2.6</b>	<b>53.4</b>		
<b>Mean</b>	<b>34.1</b>	<b>28.7</b>	<b>23.1</b>	<b>7.8</b>	<b>2.4</b>	<b>38.0</b>		
<b><i>Acacia nilotica</i></b> <b>LEAVES</b>								
<b>No. of data</b>	<b>0</b>	<b>3</b>	<b>3</b>	<b>3</b>	<b>3</b>	<b>3</b>		
<b>Low</b>		<b>11.9</b>	<b>11.3</b>	<b>5.5</b>	<b>2.0</b>	<b>42.9</b>		
<b>High</b>		<b>16.2</b>	<b>31.6</b>	<b>7.3</b>	<b>12.6</b>	<b>59.1</b>		
<b>Mean</b>		<b>13.7</b>	<b>21.4</b>	<b>6.4</b>	<b>5.6</b>	<b>52.9</b>		
<b><i>Acacia nilotica</i></b> <b>FRUIT</b>								
<b>No. of data</b>	<b>1</b>	<b>7</b>	<b>6</b>	<b>7</b>	<b>6</b>	<b>6</b>	<b>1</b>	<b>1</b>

<b>Low</b>	<b>90.1</b>	<b>10.4</b>	<b>12.3</b>	<b>4.7</b>	<b>0.8</b>	<b>51.2</b>	<b>31.6</b>	<b>22.5</b>
<b>High</b>	<b>90.1</b>	<b>13.1</b>	<b>28.5</b>	<b>13.8</b>	<b>3.0</b>	<b>68.6</b>	<b>31.6</b>	<b>22.5</b>
<b>Mean</b>	<b>90.1</b>	<b>12.0</b>	<b>17.0</b>	<b>6.6</b>	<b>2.0</b>	<b>62.3</b>	<b>31.6</b>	<b>22.5</b>
<b><i>Acacia nilotica</i></b> <b>PODS</b>								
<b>No. of data</b>	<b>0</b>	<b>1</b>	<b>0</b>	<b>1</b>	<b>0</b>	<b>0</b>	<b>1</b>	<b>1</b>
<b>Mean</b>		<b>9.7</b>		<b>6.5</b>			<b>29.8</b>	<b>22.8</b>
<b><i>Acacia nilotica</i></b> <b>SEEDS</b>								
<b>No. of data</b>	<b>0</b>	<b>1</b>	<b>0</b>	<b>1</b>	<b>0</b>			
<b>Mean</b>		<b>21.1</b>		<b>6.9</b>				
<b><i>Acacia oerfota</i></b> <b>LEAVES</b>								
<b>No. of data</b>	<b>0</b>	<b>2</b>	<b>2</b>	<b>2</b>	<b>2</b>			
<b>Low</b>		<b>32.4</b>	<b>15.8</b>	<b>8.8</b>	<b>1.7</b>	<b>41.2</b>		
<b>High</b>		<b>32.5</b>	<b>15.8</b>	<b>8.9</b>	<b>1.7</b>	<b>41.3</b>		

<b>Mean</b> <b><i>Acacia Senegal</i></b> <b>LEAVES</b>		<b>32.5</b>	<b>15.8</b>	<b>8.8</b>	<b>1.7</b>	<b>41.3</b>		
<b>No. of data</b>	<b>1</b>	<b>4</b>	<b>4</b>	<b>4</b>	<b>4</b>	<b>4</b>	<b>0</b>	
<b>Low</b>	<b>88.4</b>	<b>18.2</b>	<b>11.2</b>	<b>6.8</b>	<b>2.2</b>	<b>41.3</b>		
<b>High</b>	<b>88.4</b>	<b>20.5</b>	<b>27.7</b>	<b>8.4</b>	<b>6.7</b>	<b>55.5</b>		
<b>Mean</b>	<b>88.4</b>	<b>19.8</b>	<b>23.6</b>	<b>7.7</b>	<b>3.5</b>	<b>45.3</b>		
<b><i>Acacia Senegal</i></b> <b>FRUIT</b>								
<b>No. of data</b>	<b>0</b>	<b>2</b>	<b>2</b>	<b>2</b>	<b>2</b>	<b>2</b>	<b>0</b>	<b>0</b>
<b>Low</b>		<b>19.6</b>	<b>19.0</b>	<b>5.3</b>	<b>1.0</b>	<b>30.9</b>		
<b>High</b>		<b>22.0</b>	<b>29.6</b>	<b>7.1</b>	<b>2.1</b>	<b>43.3</b>		
<b>Mean</b>		<b>20.8</b>	<b>24.3</b>	<b>6.2</b>	<b>1.6</b>	<b>37.1</b>		
<b><i>Acacia Senegal</i></b> <b>SEEDS</b>								
<b>No. of data</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>2</b>	<b>2</b>	<b>2</b>	<b>0</b>	<b>0</b>

<b>Low</b>	<b>86.7</b>	<b>38.9</b>	<b>13.7</b>	<b>5.0</b>	<b>5.8</b>	<b>25.5</b>		
<b>High</b>	<b>86.7</b>	<b>40.1</b>	<b>20.6</b>	<b>5.9</b>	<b>10.0</b>	<b>35.4</b>		
<b>Mean</b>	<b>86.7</b>	<b>39.5</b>	<b>17.1</b>	<b>5.4</b>	<b>7.9</b>	<b>30.5</b>		
<b><i>Acacia seyal</i> FRUIT</b>								
<b>No. of data</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>2</b>	<b>2</b>	<b>2</b>	<b>0</b>	<b>0</b>
<b>Low</b>	<b>95.3</b>	<b>20.9</b>	<b>20.2</b>	<b>5.6</b>	<b>1.4</b>	<b>43.7</b>		
<b>High</b>	<b>95.3</b>	<b>21.9</b>	<b>27.4</b>	<b>9.3</b>	<b>1.9</b>	<b>47.7</b>		
<b>Mean</b>	<b>95.3</b>	<b>21.4</b>	<b>23.8</b>	<b>7.4</b>	<b>1.6</b>	<b>45.7</b>		
<b><i>Acacia sieberiana</i> LEAVES</b>								
<b>No. of data</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>0</b>	<b>0</b>
<b>Mean</b>	<b>48.8</b>	<b>15.8</b>	<b>23.6</b>	<b>11.6</b>	<b>6.3</b>	<b>42.7</b>		
<b><i>Acacia sieberiana</i></b>								

<b>FRUIT</b>								
<b>No. of data</b>	<b>2</b>	<b>4</b>	<b>3</b>	<b>4</b>	<b>3</b>	<b>3</b>	<b>1</b>	<b>1</b>
<b>Low</b>	<b>89.8</b>	<b>8.4</b>	<b>20.2</b>	<b>4.0</b>	<b>1.3</b>	<b>42.7</b>	<b>37.0</b>	<b>28.2</b>
<b>High</b>	<b>94.5</b>	<b>12.8</b>	<b>40.2</b>	<b>9.0</b>	<b>1.9</b>	<b>57.9</b>	<b>37.0</b>	<b>28.2</b>
<b>Mean</b>	<b>92.2</b>	<b>11.4</b>	<b>28.5</b>	<b>6.2</b>	<b>1.5</b>	<b>52.4</b>	<b>37.0</b>	<b>28.2</b>
<b><i>Acacia sieberiana</i></b>								
<b>PODS</b>								
<b>No. of data</b>	<b>0</b>	<b>1</b>	<b>0</b>	<b>1</b>	<b>0</b>	<b>0</b>	<b>1</b>	<b>1</b>
<b>Mean</b>		<b>9.8</b>		<b>6.8</b>			<b>40.5</b>	<b>32.7</b>
<b><i>Acacia sieberiana</i></b>								
<b>SEEDS</b>								
<b>No. of data</b>	<b>0</b>	<b>1</b>	<b>0</b>	<b>1</b>	<b>0</b>	<b>0</b>	<b>1</b>	<b>1</b>
<b>Mean</b>		<b>19.6</b>		<b>4.8</b>			<b>39.3</b>	<b>26.3</b>
<b><i>Acacia tortilis</i></b>								



<b>LEAVES</b>								
<b>No. of data</b>	<b>1</b>	<b>3</b>	<b>3</b>	<b>3</b>	<b>3</b>	<b>3</b>	<b>0</b>	<b>0</b>
<b>Low</b>	<b>90.9</b>	<b>6.5</b>	<b>9.4</b>	<b>8.7</b>	<b>2.8</b>	<b>47.8</b>		
<b>High</b>	<b>90.9</b>	<b>19.2</b>	<b>34.1</b>	<b>9.6</b>	<b>8.3</b>	<b>59.4</b>		
<b>Mean</b>	<b>90.9</b>	<b>13.0</b>	<b>18.4</b>	<b>9.0</b>	<b>5.7</b>	<b>53.9</b>		
<b><i>Acacia tortilis</i></b> <b>FRUIT</b>								
<b>No. of data</b>	<b>0</b>	<b>5</b>	<b>4</b>	<b>5</b>	<b>4</b>	<b>4</b>	<b>1</b>	<b>1</b>
<b>Low</b>		<b>12.3</b>	<b>17.5</b>	<b>4.7</b>	<b>1.7</b>	<b>49.1</b>	<b>32.4</b>	<b>24.2</b>
<b>High</b>		<b>17.8</b>	<b>24.8</b>	<b>8.4</b>	<b>3.1</b>	<b>57.9</b>	<b>32.4</b>	<b>24.2</b>
<b>Mean</b>		<b>15.8</b>	<b>20.5</b>	<b>6.5</b>	<b>2.1</b>	<b>54.1</b>	<b>32.4</b>	<b>24.2</b>
<b><i>Acacia tortilis</i></b> <b>PODS</b>								
<b>No. of data</b>	<b>0</b>	<b>1</b>	<b>0</b>	<b>1</b>	<b>0</b>	<b>0</b>	<b>1</b>	<b>1</b>
<b>Mean</b>		<b>7.8</b>		<b>6.9</b>			<b>39.4</b>	<b>26.4</b>

<b><i>Acacia tortilis</i></b>								
<b>SEEDS</b>								
<b>No. of data</b>	<b>0</b>	<b>2</b>	<b>1</b>	<b>2</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>1</b>
<b>Low</b>		<b>19.1</b>	<b>10.9</b>	<b>4.6</b>	<b>6.0</b>	<b>39.7</b>	<b>33.7</b>	<b>23.9</b>
<b>High</b>		<b>37.8</b>	<b>10.9</b>	<b>5.9</b>	<b>6.0</b>	<b>39.7</b>	<b>33.7</b>	<b>23.0</b>
<b>Mean</b>		<b>28.5</b>	<b>10.9</b>	<b>5.2</b>	<b>6.0</b>	<b>39.7</b>	<b>33.7</b>	<b>23.9</b>

**Sources: Tanner *et al.*, (1990); Le Houerou, (1980); Goodchild and McMeniman, (1987); Gohl, (1981) Skerman *et al.*, (1988).**


**Notes: NFE - nitrogen free extract**












**NDF - neutral detergent fibre**

**ADF - acid detergent fibre**



Use of Trees by Livestock: *ACACIA*



-  **Foreword**
-  **Genus *Acacia***
-  **Summary**
-  **Description and distribution**
-  **Fodder characteristics**
-  **Anti-nutritive factors**
-  **Management**
-  **Alternative uses**
-  **References and further reading**
-  **Appendix 1 - Proximate and fibre analyses of bipinnate *Acacia* spp.**
-  **Appendix 2 - Proximate analyses of *Acacia* phyllodes**



---

## Appendix 2. Proximate analyses of *Acacia phyllodes*

	Crude			Ether	
--	-------	--	--	-------	--

D:/cd3wddvd/NoExe/.../meister10.htm 75/78

	Crude protein	Crude fibre	Ash	Crude extract	NFE
<b><i>Acacia aneura</i></b>					
No. of data	4	4	4	4	4
Low	9.2	27.9	5.0	1.1	47.2
High	13.2	32.9	6.9	4.6	51.5
<b>Mean</b>	<b>11.7</b>	<b>30.0</b>	<b>5.9</b>	<b>2.7</b>	<b>49.7</b>
<b><i>Acacia cambagei</i></b>					
No. of data	1	1	1	1	1
<b>Mean</b>	<b>13.3</b>	<b>15.9</b>	<b>11.0</b>	<b>3.1</b>	<b>55.2</b>
<b><i>Acacia holosericea</i></b>					
No. of data	2	2	2	2	2
	7.8	38.6	6.0	4.0	32.3

Low	14.1	38.9	6.3	8.7	43.3
High	<b>11.0</b>	<b>38.8</b>	<b>6.2</b>	<b>6.3</b>	<b>37.8</b>
<b>Mean</b>					
<b><i>Acacia salicina</i></b>					
No. of data	1	1	1	1	1
<b>Mean</b>	<b>18.1</b>	<b>23.8</b>	<b>14.7</b>	6.0	37.4
<b><i>Acacia trachycarpa</i></b>					
No. of data	1	1	1	1	1
<b>Mean</b>	<b>14.6</b>	<b>32.6</b>	<b>7.2</b>	<b>5.0</b>	<b>40.0</b>
<b><i>Acacia tumida</i></b>					
No. of data	1	1	1	1	1

---

<b>Mean</b>	<b>15.7</b>	<b>30.2</b>	<b>7.2</b>	<b>6.7</b>	<b>40.2</b>
-------------	-------------	-------------	------------	------------	-------------

---

**Sources: Skerman *et al.* (1988); Gohl (1980); Hamel (1981).  
Notes: NFE - nitrogen free extract.**

