

light, at an hundred foot distance, and that at an hundred and twenty foot distance I could discern some of the words. When I made this tryal, its Aperture (defined next the Eye) was equivalent to more than an inch and a third part of the Object-metall. This may be of some use to those that shall endeavour any thing in *Reflexions*; for hereby they will in some measure be enabled to judge of the goodness of their Instruments, &c.

N. B. The Reader may expect in the *next Month* another Letter, which came somewhat too late to be here inserted; containing a *Table*, calculated by the same Mr. *Newton*, about the several *Apertures* and *Charges* answering the several *Lengths* of these Telescopes.

E P I T O M E

Binæ Methodi Tangentium Doctoris Johannis Wallisi Geom. Prof. Saviliani Oxoniæ; aliàs fusius & explicatius ab ipso traditæ, hîc verò ob angustiam loci compendifactæ: In quarum Schematismis si forsan literæ quædam redundaverint, illæ ad ea pertinere censendæ sunt, quæ in ampliori ejusdem Scripto continentur, hîc vero dictâ de causa omittantur.

HAbes hic (*Clarissime vir*) eorum summam (*strictim traditam*) quæ fusius scripseram, meas de Tangentibus Methodos spectantia, duas potissimum quibus præsertim utor; alteram in Speciebus, alteram in Lineis; utramque generali formâ facile explicablem.

Priorem adhibeo Con.Se&. prop. 23, 30, 36, 46, 49. & passim alibi. Quæ hæc est.

Expositâ Curvâ Aa, (put à Parabola, fig. 4.) quam in a tangat a F, diametro VDA occurrens in F; ordinatum applicentur a V, & DOT curvæ in O & tangentî in T occurrens. Ponatur autem Va=b, VA=v, VF=f, VD=a, adeoque DA=v+a, DF=f+a;

Est (propter similia triangula) VF.DF :: Va.DT = $\frac{f+a}{f} b$.

Item, si tangens sit ultra curvam, DT > DO; si citra, DT < DO: Nempe, DT=DO si intelligatur D in V; sed, si extra V, DT vel DO major prout tangens est ultra citrave curvam.

Tum,



Tum, habita ipsius DO designatione quæ sit expositæ curvæ accommodata; (puta, in Parabolæ, propter AV . $AD :: Vaq.$. $DOq = \frac{v^{\pm a}}{v} b^2$; $DO = b \sqrt{\frac{v^{\pm a}}{v}}$:) fiat debita reduc[t]io, (puta, propter $\frac{f^{\pm a}}{f} b > b \sqrt{\frac{v^{\pm a}}{a}}$, adeoque $\frac{f^2 \pm 2fa + a^2}{f^2} > \frac{v^{\pm a}}{v}$, & $f^2 v \pm 2fv a + va^2 > f^2 v \pm f^2 a$; deletis utrinque æqualibus, hoc est, iis omnibus in quibus a non conspicitur; cæteris que per \pm a divisis: $2fv \pm va > f^2$.)

Tandem (qui methodi nucleus est) posito D in V , (quod sit $a=0$, adeoque evanescent ipsius multiplæ omnia,) æquatio exhibebit f quæsitam (puta $2fv \pm va = 2fv = f^2$, adeoque $2v = f$.)

Hanc (locis citatis) accommodatam videas Parabolæ, Ellip[s]i, Circuleve, Hyperbolæ, Paraboloidibus omnibus, (quibus & harum Reciprocas accenseo,) atque alibi aliis.

Cissoidi (fig. 5.) sic accommodes. Est (per cap. 5. pr. 29. de Motu) $Va = b = \frac{v^2}{v+h}$, (posito s pro sinu recto in circulo generante, cuius radius r, sinus versus v, & $2r \cdot v = h$, & $h \cdot v = 2a$,) adeoque (substitutis $v \pm a$ pro v, & $h \mp a$ pro h,) $\frac{v^2 \pm 2va + a^2}{v(v+h) \pm 2xa - a^2} (= DO) > (DT = \frac{f^{\pm a}}{f} b =) \frac{f^{\pm a}}{f} \times \frac{v^2}{v+h}$. Ergo (sumptis quadratis, & multiplicando decussatim,) $f^2 v^5 h + 6f^2 v^3 ha^2 + f^2 vha^4 \pm 4f^2 v^4 ha \pm 4f^2 v^2 ha^3 > f^2 v^5 h \pm 2fv^3 ha \pm v^5 ha^2 \pm 2f^2 v^4 xa \pm 4f^2 v^4 xa^2 \pm 2v^4 xa^3 - f^2 v^4 a^2 \mp 2fv^4 a^3 - v^4 a^4$: item (deletis utrinque æqualibus, cæteris que per \pm va divisis) $\pm 6f^2 v^2 ha \pm f^2 ha^3 + 4f^2 v^3 h + 4f^2 vha^2 > 2fv^4 h \pm v^4 ha \pm 2f^2 v^3 xa \pm 4f^2 v^3 xa \pm 2v^3 xa^2 \mp f^2 v^3 a^2 - 2fv^3 a^2 \mp v^3 a^3$. Denique (posito D in V , quo evanescat a cum suis multiplis, cæteris que per \pm fv^3 divisis) fiet æquatio $2fh - fa = vh$, adeoque $\frac{vh - 5a^2}{2h - x - \pm h - 3r - v} = f$.

Idem succedet, sumptâ, pro VA , diametro TA , (cui tangens occurrat in Φ) aliâ Ψ . Item, si exponeretur curva quæ ordinatas non habeat, sed quæ his equipollent; ut sunt, in Spirali, crescentes radii.

Sed & calculi magna pars præverti potest; omissis ab initio (ut pote post rei sciendis) terminis iis ubi habetur x^2 vel superior hujus potestas; item iis in quibus nec a conspicitur, nec sunt in a ducendi, (ut pote æqualibus utrinque proditur.) Exempli gratiâ,

In Conchoide, (fig 6.) cuius ordinata VMa constat ex sinu recto $VM=s=\sqrt{vh}$, & tangente $Ma=CH=\frac{s}{x}r$, (si sit $CP=CA=r$, adeoque $CH=AS$;) saltem $=\frac{s}{x}r$ (posito $CP=s$;) adeoque $Va=b=s+\frac{s}{x}r=\frac{x+r}{x}s=\frac{h}{x}s$, saltem $\frac{x+r}{x}s=\frac{s}{x}s=\frac{s}{x}\sqrt{vh}$ (posito $x+s=n$) Ergo $DT=\frac{f \pm a}{a}b=\frac{f \pm a}{f_n n}\sqrt{vh} \geq DO=\frac{n^2 f^2}{x^2 + a^2} \sqrt{vh + 2xa - :}$ (omittit a^2 , quia post delendum, indeque oriunda, & sic semper:) &, sumptis quadratis, $\frac{f^2 + 2fa + a^2}{f^2 x^2} n^2 v h \geq \frac{n^2 v h + 2na^2 - 2va^2}{x^2 + 2xa - :} v h a^2$ (hoc est, \geq supra, sed \leq infra, punctum flexus contrarii.) Et, decussatim multiplicando; omissis (ut precipitur) $f^2 x^2 n^2 v h$ utrobique, omnibusque a^2 multipliis; ceterisque per $\pm a$ divisis; $2f_n^2 v h a^2 - 2f^2 n^2 v h a^2 \geq 2f^2 n^2 x^3 - 2f^2 n v h a^2$: adeoque (posito D in V ,) $n v h a^2 = f_n v h + f_n x^2 - f v h a^2 = f_n r^2 - f v h a^2$ (propter $v h + x^2 = s^2 + x^2 = r^2$), & $f = \frac{\sqrt{v h + r^2}}{n^2 + x^2 - 2f^2} n r$. Et quidem, in primaria, (propter $h = u$), $f = \frac{s^2 x}{x^2 - s^2}$.

In Figura Tangentium (fig.7.) quæ à Conchoide differt, excepto quadrante genitore; idem erit processus, nisi quod, propter $Va=Ma=\frac{p}{n}s$ (non $\frac{n}{x}s$), prodibit (sive in primaria, sive in protracta contractave,) $f = \frac{\sqrt{v h + s^2}}{r^2} x$.

In Figura Secantium (fig.8.) propter $Va=b=\frac{r^2}{x}$; erit $DO = \frac{r^2}{x^2 + a^2} \geq \frac{f \pm a}{f_n} r^2 = DT$. adeoque $f = x$.

Cumque hæc curva sit Hyperbola (per pr.30.cap.5. & pr.1.cap.15. de Motu,) cuius Asymptotæ CA , CB : eadem tangens habetur per pr. 36.Con. sect. Cumque ordinata ad asymptotas (per pr. 94, 95, Arith. Infin.) sint series Reciproca Primanorum (quæ ad Paraboloidium genus spectat, verticem habens C, exponentem -1 ,) habetur eadem tangens per prop. 49. Con. Sect. (eademque est expedita methodus pro hyperbolæ cuiusvis tangente per asymptotam inveniendâ.) Quippe, in Paraboloidibus omnibus, ut intercepta diameter VC , ad VF , sic 1 ad exponentem: hoc est, in praesenti casu, ut 1 ad -1 ; adeoque $VC=VF$, sed (propter contraria signa $+$ -) ad contrarias partes.

Notandum hic; in Parabolâ, Paraboloidë, Hyperbolâ, Ellipſe, &c. figurâve Sinum (rectorum, versorumve,) Arcuum, Tangentium,

gentium, Secantium, &c. aliâve cuius constructio est similaris; protractio contractiove figuræ (seu mutatio Lateris recti, aut quod bujus instar est,) non mutat punctum F, (eo quod Latus-rectum aequationem quæ longitudinem VF determinat non ingrediatur, utut eam ingrediatur quæ determinat longitudinem Va, mutetque angulos ad a & F:) sed ubi constructio est Dissimilaris, ut in Cycloïde & Conchoïde (propter ordinatam illic ex Sinu & Arcu, hic ex Sinu & Tangente, conflatam,) aliisque istiusmodi, res securus est: id quod una pars (ut Arcus in Cycloïde & Tangens in Conchoïde) protractitur contrahiturve, manente alterâ (putâ, in utrisque Sins recto) ut in primaria.

Idemque dicendum de Angulo applicationis (ad V,) cuius mutatio non mutat longitudinem VF, sed neque Va, quia neutrius ingreditur equationem. Atque hinc fit, quod in figura Scalena, quæ ordinatas contrarias, utrinque ad V positas, spectant tangentes, utut inæquales, in eodem F convenient. Sed & (ut hoc obiter moneam) quadratorum aggregatum habent idem atque in erectâ; nempe semper = 2 Va q + 2 VF q.

Estque hæc mihi methodus pro Maximis & Minimis in omne genus quantitatibus.

Methodus altera (sesundum tradita de Angulo Contactus & Arithm. Infin.) curvam considerat tanquam ex pariculis conflatam infinitè exiguis, sed certam positionem habentibus; eandem nempe (propter contactus angulum sive nullius magnitudinis sive infinitè exigua) cum recta ibidem tangente: adeoque cum hac (respectu cuiusvis rectæ) pariter declivem, (ut est Montis Aa fig. 4, 5, declivitas in a, eadem quæ tangentis a F.) Cujus ergo quæque particula (per cap. 2. de Motu) est in eâ ratione magis longa (quam est respectiva expositæ rectæ particula æquè-alta) quâ est minus declivis; puti a T quam VD: Unde, propter mutatam in singulis punctis declivitatem, oritur series longitudinum inæqualium in curvâ, seriei æqualium in rectâ, respondens; curvæ ad rectam rationem exhibens. Atque hinc methodus mea pro curvis rectificandis, (schol. prop. 38. Ar. Infin. insinuata,) quam prosequor tractatu de Eucl. item de motu cap. 5. prop. 13. & seqq. Cujus aliqua pars est hæc de Tangentibus, ut quæ non totam declivitatem seriem perpendit, sed eam quæ est in extremitate puncto.

Hanc respectivam particularam longitudinem, alias insinuatum eunt

eunt (motu forinsecus assumpto) per motuum quibus transfigantur $\text{io}\chi\text{gōvōw}$ celeritatem. (Quippe idem est, in Motu, Celeritas, atque hæc, in Situ (propter positionem obliquam seu minus declivem) respectiva Longitudo.) Aptissimè quidem in lineis a motu primitus oriundis, (puta, Cycloide, Conchoide, Spirali, Quadratice, &c.) nec ineptè in aliis, quæ fangi saltem possunt istiusmodi motibus describi.

Præsumo autem (ex prop. 15. cap. 2. de Motu) eam esse curvæ in quovis puncto directionem, adeoque & declivitatem, quæ est rectæ ibidem tangentis: Item (ex prop. 6. cap. 10.) Motus compositi directionem esse in Diagonio parallelogrammi, cuius latera & anguli exhibeant componentium celeritates & directiones.

Intelligatur jam (fig. 4.) Aa parabola, describi motu composito, ex æquabili secundum AY vel V_a , cuius itaque particulae $\text{io}\chi\text{gōvōw}$ (per pr. 3. cap. 10. de Motu) sunt series Primæorum, quæ ad se rem totidem ultimæ æqualium, (hoc est, ad rectam $\text{io}\chi\text{gōvōw}$ celeritate in a acquisitam transfigendam,) est ut 1 ad 2, (per Ar. Infin. pr. 64. vel pr. 1. cap. 5. de Motu.) Adeoque, sumpta $VF = 2VA$, & compagno $FVaw$ parallelogrammo; juncta aF est Tangens.

Idem similiter obtinebitur in Paraboloidibus quibuscunque, ope prop. 2, 5, 6, 7, de Motu.

Atque inde facile (vel ex iisdem principiis) ostenditur; si intel ligatur Fig. AYv sic constituta, ut momenta (respectu AF) ordinatum Yv , y_w , sint ipsis Ya , yO , ordinatis proportionalia; erunt Celeritates acquisitæ in a , o , seu V , D , (posita AY linea motus æquabilis) rectis $Y_{\frac{a}{2}}$, y_w , proportionales: Et consequenter, ut AuY (illarum aggregatum) ad $AFuY$ (aggregatum totidem maximæ æqualium,) sic VA (aggregatum celeritatum seu particularum crescentium) ad (aggregatum totidem maximæ æqualium) TF .

Spiralis ASa (fig. 9.) punctum a designatur motu composito ex recto per Aa , & circulari per Va , æquabilibus utrisque & $\text{io}\chi\text{gōvōw}$. Ergo, sumpta circuli tangente $w=aV$, & completo $AawF$ parallelogrammo; juncta aF Spiralem tanyet.

Vnde statim emergit Archimedea quadratura (sive Circuli sive Sectoris cuiusvis) propter $AF = aw = aV$.

Sin motuum alter, puta Aa , sit acceleratus vel retardatus; pro

αA , sumenda erit αB (in ea ad illuminatione quam illa postulat acceleratio seu retardatio,) eritque diagonum $\alpha\beta$, Tangens quæfita.

Quadratricis $A\alpha B$ (fig. 10.) punctum α designatur motu composito ex recto per $v\alpha$, & circulari in $\Upsilon\alpha$ (æquabilibus & in æquibus.) Ergo, sumpta tangente $\alpha V=\alpha\Upsilon$, & completo parallelogrammo $V\alpha\omega F$, juncta αF tanget Quadratricem.

Atque hinc alia quadratura, per Tangentem quadratricis, propter $vF=\alpha V=\alpha\Upsilon$.

IIIa per quadratricis Basin, sic elicetur. Positis $CA=r$, $AQ=q$, $\Upsilon Z=x$, $QR=a$. Erit (propter Quadratricis constructionem) AQ . $RQ :: AC.\alpha AE = . \frac{1}{q} r :: \Upsilon Z. \alpha Z = \frac{1}{q} x$. Estque $\alpha Z > \alpha E$ sumpto ubi vis in AB punto α , præterquam in B , quo casu (evanescente utraque) erit $\alpha Z=\alpha E$, adeoque $x=r$; hoc est, $\Upsilon Z=XB=AC$. Sed & vE communis tangens utriusque curvæ XB , AB .

Cycloïdis (fig. 11.) punctum α describitur motu composito, ex recto in αV , & circulari in $\alpha\beta$ (æquabilibus & æquè-velocibus.) Ergo, sumpta tangente $\alpha\omega=\alpha V$, & completo $V\alpha\omega F$ parallelogrammo, juncta αF Cycloidem tanget. Et quidem, propter Ang. $v\alpha F$ ($=\alpha\beta F = \frac{1}{2} \alpha CF$) $= \frac{1}{2} v\alpha V$, occurret circuli $\alpha\beta$ erectæ diametro in vertice.

In secundariis (contractâ protractâve) sumenda erit $\alpha\omega$ ad αV , in ea ratione major minore, quâ est celeritas motûs circularis ad celeritatem recti.

In Figura Arcuum, Sinuumve, (fig. 12.) procedendum ut in Cycloide, nisi quod (propter exemptum semicirculum genitorem) pro tangentे $\alpha\omega$ illic (quæ hic est at) sumenda erit erecta $\alpha\omega$ æqua-alta.

Conchoidis (fig. 6.) punctum α designatur motu composito, ex æquabili circulari in $\alpha\beta$ (hujusve tangentे $\alpha\omega$) & recto in $\alpha\Upsilon$ acelerato pro incremento tangentium: que quidem acceleratio duplex est, altera propter declivitatis angulum $\beta\alpha\Upsilon$, hoc est, $v\alpha\Upsilon$, continuè crescentem; altera propter radii in secantem protractionem, continuè item crescentem. Propter priorem, ducta tangentē $\alpha\omega$ (que occurrat in v regularē CH .) recta $v\zeta$ (parallela recte $PH\omega$.) occurrat $\alpha\Upsilon$ in ζ : Propter posteriorēm, eidem $v\zeta$ protracta occurrat tangentē verticis in Z : indeque $Z\Upsilon$ restat $v\alpha X$ parallela; adeoque $\alpha\Upsilon=XZ$.

$\alpha \ell :: CM. \alpha \ell S :: P\mu. PH.$ Completo denique τ auf parallelogrammo, juncta αf tanget conchoidem.

In secundariis (ubi non est $CP=CA,$) sumenda erit $\alpha \tau$ ad jam designatam, ut est CP ad $CA.$

In Figura Tangentium (fig. 7.) propter exemptum Conchoidi quadrantem genitorem, pro tangentे αv illio (quæ hic est αr) sumenda erit erecta αv æquè-alta.

Pluribus exemplis preferendis supersedeo. Moneo tamen, utramvis Methodum, utut tangentibus rectis hic accommodatam, extendere posse ad mutuum Curvarum tactum. Puta; si, pro $FV\alpha$ triangulo (fig. 4,5,) intelligatur Hyperbola; recta DT , quæ hic insinuitur charactere qui triangulo conveniet. subire tum debebit characterem Hyperboles; cujus vertex F simili processu queratur. Similisque in posteriori methodo accommodatus est linearum ductus. Et quidem, cum curvam $A\alpha$ tangens recta αF , sit etiam tangens communis curvarum omnium, expositam ibidem tangentium; prout hic, ex data $A\alpha$ curva queritur recta αF , sic ex hac datâ (per eandem methodum inversam) querenda erit alia tangens curva, modo satis sit determinata.

Sed ampliandum non est. Tu itaque Vale.

Tuus

Oxonie abe 15.

Febr. 1671.

Johannes Wallis.

Extract

