

Complex Functions (1A)

Copyright (c) 2011 Young W. Lim.

Permission is granted to copy, distribute and/or modify this document under the terms of the GNU Free Documentation License, Version 1.2 or any later version published by the Free Software Foundation; with no Invariant Sections, no Front-Cover Texts, and no Back-Cover Texts. A copy of the license is included in the section entitled "GNU Free Documentation License".

Please send corrections (or suggestions) to youngwlim@hotmail.com.

This document was produced by using OpenOffice and Octave.

Derivatives

the complex function f is defined in a neighborhood of a point z_0


Derivative of f at z_0

$$f'(z) = \frac{df}{dz} = \lim_{\Delta z \rightarrow 0} \frac{\Delta f}{\Delta z} = \lim_{\Delta z \rightarrow 0} \frac{f(z_0 + \Delta z) - f(z_0)}{\Delta z}$$

provided that this limit **exists**

f is said to be **differentiable** at z_0

Δz can approach zero
from any convenient direction



Analyticity

f **differentiable** at z_0
 f **differentiable** at every point
in some *neighborhood* of z_0



the complex function f is
analytic at a point z_0

A complex function can be
differentiable at a point z_0
but differentiable nowhere else

$$f(z) = z\bar{z} = |z|^2 = x^2 + y^2$$

$$\lim_{\Delta z \rightarrow 0} \frac{(x+\Delta x)^2 + (y+\Delta y)^2 - x^2 - y^2}{\Delta x + i\Delta y}$$

A function that is **analytic**
at **every** point z
:analytic function

$$f(z) = z^2$$

Analytic Functions

$$f'(z) = \frac{df}{dz} = \lim_{\Delta z \rightarrow 0} \frac{\Delta f}{\Delta z}$$

$$\Delta f = f(z + \Delta z) - f(z)$$

$$\Delta z = \Delta x + i\Delta y$$

$f(z)$: **analytic** in a region



$f(z)$ has a (unique) derivative at every point of the region

$f(z)$: **analytic** at a point
 $z = a$



$f(z)$ has a (unique) derivative at every point of some small circle about $z = a$

Singular Point

Regular point of $f(z)$



a point at which
 $f(z)$ is **analytic**

Singular point of $f(z)$



a point at which
 $f(z)$ is **not analytic**

Isolated Singular point of $f(z)$



a point at which
 $f(z)$ is **analytic**
everywhere else
inside some small circle
about the singular point

Cauchy-Riemann Condition (1)

$$f(z) = u(x, y) + iv(x, y)$$

: **differentiable** at a point

$$z = x + iy$$



$$\frac{\partial u}{\partial x} = + \frac{\partial v}{\partial y}$$

$$\frac{\partial v}{\partial x} = - \frac{\partial u}{\partial y}$$

**Necessary Condition
for Analyticity**

$$f(z) = u(x, y) + iv(x, y)$$

$$\frac{\partial}{\partial x}$$

$$\frac{\partial}{\partial y}$$

$$f(z) = u(x, y) + iv(x, y)$$

$$\frac{\partial}{\partial x} \quad \frac{\partial}{\partial y}$$

Cauchy-Riemann Condition (2)

$f(z) = u(x, y) + iv(x, y)$: **differentiable** at a point $z = x + iy$

➡ $f'(z)$ exists


$$f'(z) = \lim_{\Delta z \rightarrow 0} \frac{f(z_0 + \Delta z) - f(z_0)}{\Delta z} \quad \Delta z = \Delta x + i\Delta y$$

$$= \lim_{\Delta z \rightarrow 0} \frac{u(x+\Delta x, y+\Delta y) + iv(x+\Delta x, y+\Delta y) - u(x, y) - iv(x, y)}{\Delta z}$$

horizontal approach ➡ $\Delta x \rightarrow 0$
 $\Delta z \rightarrow 0$ $\Delta y = 0$

vertical approach ➡ $\Delta y \rightarrow 0$
 $\Delta z \rightarrow 0$ $\Delta x = 0$


must have
the same
 $f'(z)$



Cauchy-Riemann Condition (3)


horizontal approach $\Delta z \rightarrow 0 \Rightarrow \Delta x \rightarrow 0$ $\Delta y = 0$

$$\begin{aligned} f'(z) &= \lim_{\Delta z \rightarrow 0} \frac{u(x+\Delta x, y+\Delta y) + i v(x+\Delta x, y+\Delta y) - u(x, y) - i v(x, y)}{\Delta z} \\ &= \lim_{\Delta z \rightarrow 0} \frac{u(x+\Delta x, y) - u(x, y)}{\Delta x} + i \frac{v(x+\Delta x, y) - v(x, y)}{\Delta x} \\ &= \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} = \frac{\partial f}{\partial x} \end{aligned}$$



vertical approach $\Delta z \rightarrow 0 \Rightarrow \Delta y \rightarrow 0$ $\Delta x = 0$


$$\begin{aligned} f'(z) &= \lim_{\Delta z \rightarrow 0} \frac{u(x+\Delta x, y+\Delta y) + i v(x+\Delta x, y+\Delta y) - u(x, y) - i v(x, y)}{\Delta z} \\ &= \lim_{\Delta z \rightarrow 0} \frac{u(x, y+\Delta y) - u(x, y)}{i \Delta y} + i \frac{v(x, y+\Delta y) - v(x, y)}{i \Delta y} \\ &= -i \frac{\partial u}{\partial y} + \frac{\partial v}{\partial y} = -i \frac{\partial f}{\partial y} \end{aligned}$$



Cauchy-Riemann Condition (4)


horizontal approach $\Delta z \rightarrow 0 \Rightarrow \Delta x \rightarrow 0$ $\Delta y = 0$

$$f'(z) = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} = \frac{\partial f}{\partial x}$$



vertical approach $\Delta z \rightarrow 0 \Rightarrow \Delta y \rightarrow 0$ $\Delta x = 0$

$$f'(z) = -i \frac{\partial u}{\partial y} + \frac{\partial v}{\partial y} = -i \frac{\partial f}{\partial y}$$



$$\Rightarrow f'(z) = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} = -i \frac{\partial u}{\partial y} + \frac{\partial v}{\partial y} \Rightarrow \frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} \quad \frac{\partial v}{\partial x} = -\frac{\partial u}{\partial y}$$

To Be Analytic (1)

$f(z) = u(x, y) + iv(x, y)$: **analytic** in a domain D



$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}$$

$$\frac{\partial v}{\partial x} = -\frac{\partial u}{\partial y}$$

$f(z) = u(x, y) + iv(x, y)$: **analytic** in a domain D



$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}$$

$$\frac{\partial v}{\partial x} = -\frac{\partial u}{\partial y}$$

$u(x, y), v(x, y)$: **continuous** on in a domain D


$\frac{\partial u}{\partial x}, \frac{\partial u}{\partial y}, \frac{\partial v}{\partial x}, \frac{\partial v}{\partial y}$: **continuous** on in a domain D

To Be Analytic (2)

if the real functions u and $v(x,y)$ are continuous
and have continuous first order partial derivatives
in a neighborhood of z ,
and if $u(x,y)$ and $v(x,y)$ satisfy
the Cauchy-Riemann equations at the point z ,

then the complex function $f(z) = u(x,y) + iv(x,y)$
is **differentiable** at z
and $f'(z)$ is as follows.

$$f'(z) = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} = \frac{\partial v}{\partial y} - i \frac{\partial u}{\partial y}$$



Derivatives

$f(z) = u(x, y) + iv(x, y)$: **analytic** in a region R



derivatives of all orders at points inside region


$f'(z_0), f''(z_0), f^{(3)}(z_0), f^{(4)}(z_0), f^{(5)}(z_0), \dots$



Taylor series expansion about any point z_0 inside the region

The power series converges inside the circle about z_0

This circle extends to the nearest **singular point**



Laplace Equation

$f(z) = u(x, y) + i v(x, y)$: **analytic** in a region R

➡ $u(x, y), v(x, y)$ satisfy Laplace's equation in the region
harmonic functions

$u(x, y), v(x, y)$ satisfy Laplace's equation in simply connected region

➡ Real / imaginary part of an analytic function $f(z)$

References

- [1] <http://en.wikipedia.org/>
- [2] <http://planetmath.org/>
- [3] M.L. Boas, "Mathematical Methods in the Physical Sciences"